

# Scheme to treat and prevent **Megabacteria** in birds

## To consider:

- 1.- They are classified as a fungus, although it has the form of a bacterium so big.
- 2.- It is installed in the proventriculus of the birds (the union of the crop with the gizzard) causing the necrosis of these tissues and causing diarrhea, anemia and in many cases death.
- 3.- It is very frequent that Megabacteria are accompanied by other diseases caused by bacteria.
- 4.- It is transmitted through the excrements.

## Products needed for treatment:



- **Total Disinfection Solution**, to control the spread of germs through water (one of the main routes of spread of the disease).



- **Megabac Plus MS**, to control and prevent the proliferation of bacteria associated with Megabacteria.



- **Fungi-Plus MS**, a powerful fungicide to fight Megabacteria.



- **Prowins Probibac Bird**, the perfect combination of Probiotics, Prebiotics, Electrolytes, Vitamins and Amino Acids, which guarantees a perfect intestinal flora throughout the year, even in periods of high stress. Specifically developed for Birds and Cage Birds

## Weekly schedule for the treatment and prevention of Megabacteria

Days	Products to use:
Day 1	<b>Total Disinfection Solution</b> (10ml per 1L of water) + <b>Megabac Plus</b> (4g per 1L of water)
Day 2	<b>Megabac Plus</b> (4g per 1L of water) + <b>Fungi-Plus</b> (5ml per 1L or 1kg of egg food) + <b>Probibac Bird</b> (1 tablespoon per 1kg of egg food)
Day 3	<b>Megabac Plus</b> (4g per 1L of water) + <b>Fungi-Plus</b> (5ml per 1L or 1kg of egg food) + <b>Probibac Bird</b> (1 tablespoon per 1kg of egg food)
Day 4	<b>Total Disinfection Solution</b> (10ml per 1L of water) + <b>Megabac Plus</b> (4g per 1L of water)
Day 5	<b>Megabac Plus</b> (4g per 1L of water) + <b>Fungi-Plus</b> (5ml per 1L or 1kg of egg food) + <b>Probibac Bird</b> (1 tablespoon per 1kg of egg food)
Day 6	<b>Megabac Plus</b> (4g per 1L of water) + <b>Fungi-Plus</b> (5ml per 1L or 1kg of egg food) + <b>Probibac Bird</b> (1 tablespoon per 1kg of egg food)
Day 7	<b>Total Disinfection Solution</b> (10ml per 1L of water) + <b>Megabac Plus</b> (4g per 1L of water)

✓ This scheme must be applied throughout the breeding season, ending before the change.

## Megabacteria

