

Salmonellosis Treatment Plan for Pigeons (Individual treatment)

To consider:

- 1.- Very common disease in all types of birds, which affects both chicks as adults.
- 2.- Pigeons that pass the disease can become carriers.
- 3.- It can present in several ways: Intestinal (acute greenish diarrhea), joint (wings falling and legs shrunken), organic (affecting vital organs) and nervous (loss of balance, paralysis, turns of the neck similar to those caused by paramyxovirosis)
- 4.- It spreads through food, water, by air and through the egg shell.

Products needed for treatment:



- **Total Disinfection Solution**, to control the spread of germs.



- **Baytril Master Formula**, for the elimination of the causative agent of the disease (salmonella typhimurium)



- **Probibac Caps**, to regenerate the intestinal flora and replenish all fluids lost during the disease process.

Scheme to follow for the treatment of Salmonellosis

Days	Products to use
Day 1	Total Disinfection Solution (10 ml per 1 liter of drinking water)
Day 2	Baytril Master formula (3 direct drops to the peak) + Probibac Caps (1 cap per pigeon)
Day 3	Baytril Master formula (3 direct drops to the peak) + Probibac Caps (1 cap per pigeon)
Day 4	Total Disinfection Solution (10 ml per 1 liter of drinking water)
Day 5	Baytril Master formula (3 direct drops to the peak) + Probibac Caps (1 cap per pigeon)
Day 6	Baytril Master formula (3 direct drops to the peak) + Probibac Caps (1 cap per pigeon)
Day 7	Total Disinfection Solution (10 ml per 1 liter of drinking water)
Day 8	Baytril Master formula (3 direct drops to the peak) + Probibac Caps (1 cap per pigeon)
Day 9	Probibac Caps (1 cap per pigeon)
Day 10	Probibac Caps (1 cap per pigeon)



This plan must be applied for 10 days. At the end of this period, we evaluate and if the pigeons are not yet recovered, we wait 4-6 days and repeat the treatment.

Salmonellosis

