

Treatment against Coccidiosis in Pigeons

To consider:

- 1.- Intestinal disease that affects all types of birds, affecting both chicks as adults.
- 2.- It can present acute and recognizable symptoms: watery, greenish diarrhea, with bad smell and apathetic and visibly diseased pigeons (Acute evolution of the illness)
- 3.- It can also present very mild symptoms, difficult to diagnose: seemingly healthy pigeon, with little vitality, poor performance on flights and, sometimes, with soft excrement (Asymptomatic evolution)
- 4.- Sometimes it is confused with salmonellosis, e-coli, intestinal worms.

Necessary products:



- **ESB4 Plus**, for the control and elimination of coccidiosis.



- **Prowins B12**, to strengthen the immune system and boost the effectiveness of ESB4 Plus



- **Probibac Forte**, to regenerate the intestinal flora.

Scheme to follow for the treatment of Coccidiosis

Days	Products to use:
Day 1	- In the drinking water: ESB4 Plus (5g per liter of water) + Prowins B12 (15ml per liter of water) - In the food: Probibac Forte (7,5g per kilo of food)
Day 2	- In the drinking water: ESB4 Plus (5g per liter of water) + Prowins B12 (15ml per liter of water) - In the food: Probibac Forte (7,5g per kilo of food)
Day 3	- In the drinking water: ESB4 Plus (5g per liter of water) + Prowins B12 (15ml per liter of water) - In the food: Probibac Forte (7,5g per kilo of food)
Day 4	- In the drinking water: ESB4 Plus (5g per liter of water) + Prowins B12 (15ml per liter of water) - In the food: Probibac Forte (7,5g per kilo of food)
Day 5	- In the drinking water: ESB4 Plus (5g per liter of water) + Prowins B12 (15ml per liter of water) - In the food: Probibac Forte (7,5g per kilo of food)
Day 6	- In the drinking water: ESB4 Plus (5g per liter of water) + Prowins B12 (15ml per liter of water) - In the food: Probibac Forte (7,5g per kilo of food)
Day 7	- In the drinking water: ESB4 Plus (5g per liter of water) + Prowins B12 (15ml per liter of water) - In the food: Probibac Forte (7,5g per kilo of food)



This plan must be administered for 7 consecutive days.

Coccidiosis

