

# Individual Treatment against Coccidiosis in Pigeons

## To consider:

- 1.- Intestinal disease that affects all types of birds, affecting both chicks as adults.
- 2.- It can present acute and recognizable symptoms: watery, greenish diarrhea, with bad smell and apathetic and visibly diseased pigeons (Acute evolution of the illness)
- 3.- It can also present very mild symptoms, difficult to diagnose: seemingly healthy pigeon, with little vitality, poor performance on flights and, sometimes, with soft excrement (Asymptomatic evolution)
- 4.- Sometimes it is confused with salmonellosis, e-coli, intestinal worms.

## Necessary products:



- **Baycox**, for the control and elimination of coccidiosis.



- **Probibac Caps**, to regenerate the intestinal flora and strengthen the defenses.

## Scheme to follow for the treatment of Coccidiosis

Days	Products to use:
Day 1	- In the drinking water: <b>Baycox</b> (4 ml per liter of water or 3 drops direct to the peak) - <b>Probibac Caps</b> ( 1 cap per pigeon )
Day 2	- In the drinking water: <b>Baycox</b> (4 ml per liter of water or 3 drops direct to the peak) - <b>Probibac Caps</b> ( 1 cap per pigeon )
Day 3	- In the drinking water: <b>Baycox</b> (4 ml per liter of water or 3 drops direct to the peak) - <b>Probibac Caps</b> ( 1 cap per pigeon )
Day 4	- In the drinking water: <b>Baycox</b> (4 ml per liter of water or 3 drops direct to the peak) - <b>Probibac Caps</b> ( 1 cap per pigeon )
Day 5	- In the drinking water: <b>Baycox</b> (4 ml per liter of water or 3 drops direct to the peak) - <b>Probibac Caps</b> ( 1 cap per pigeon )



This plan must be administered for 5 consecutive days.

## Coccidiosis

